

Rec'd PCT/PTO 12 JUL 2004

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
17 July 2003 (17.07.2003)

PCT

(10) International Publication Number  
**WO 03/057250 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 39/395**

(21) International Application Number: **PCT/GB03/00105**

(22) International Filing Date: 13 January 2003 (13.01.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0200657.5 12 January 2002 (12.01.2002) GB

(71) Applicant (for all designated States except US): **ANTI-SOMA PLC** [GB/GB]; West Africa House, Hanger Lane, Ealing, London W5 3QR (GB).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **ROWLINSON-BUSZA, Gail** [GB/GB]; Antisoma Research Laboratories, St. George's Hospital Medical School, Cranmer Terrace, London SW17 0QS (GB).

(74) Agent: **THOMAS, Philip, J., D.**; Eric Potter Clarkson, Park View House, 58 The Ropewalk, Nottingham NG1 5DD (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **CANCER TREATMENT**

(57) Abstract: The invention relates to a therapeutic system comprising (i) a radiolabelled antibody, which binds selectively to polymorphic epithelial mucin (PEM) such as the monoclonal antibody HMFG-1, and (ii) a chemotherapeutic agent, such as Taxotere<sup>®</sup>. The radiolabelled antibody and chemotherapeutic agent are administered in combination with one another to produce a synergistic therapeutic effect.

**BEST AVAILABLE COPY**

**WO 03/057250 A1**

**THIS PAGE BLANK (USP)**

The present invention relates to materials and methods for the treatment of cancer. In particular, the invention relates to a therapy comprising the administration of a radiolabelled antibody, which binds selectively to polymorphic epithelial mucin (PEM) in combination with a chemotherapeutic agent.

DeNardo *et al.* (April 1997) *P.N.A.S. USA* 94 pp. 4000-4004 mention that a synergistic therapeutic effect can be obtained in a mouse model of breast cancer by administering a yttrium 90-labelled chimeric L6 antibody ( $^{90}\text{Y}$ -ChL6) 6 or 24 hours before Taxol® (paclitaxel) administration. However, no synergistic effect was observed when Taxol® was administered 24-27 hours before  $^{90}\text{Y}$ -ChL6.

DeNardo *et al.* conclude as follows:  $^{90}\text{Y}$ -ChL6 and Taxol® can be given in a sequence that enhances therapeutic efficacy. Over time, after injection,  $^{90}\text{Y}$ -ChL6 binds to malignant cells as it circulates and unbound  $^{90}\text{Y}$ -ChL6 is cleared from normal tissues. Thus a "window" in time exists when there is ongoing tumour irradiation but little concurrent normal tissue irradiation. Given in this window, Taxol®, a small molecule rapidly taken up by the tumour, enhances the therapeutic effect of  $^{90}\text{Y}$ -ChL6 on targeted malignant cells. The optimum time for Taxol® administration is 6-24 hours after  $^{90}\text{Y}$ -ChL6.

ChL6 consists of a human IgG constant region and the Fab' region of murine monoclonal antibody (mAb) L6. ChL6 reacts with an integral membrane glycoprotein expressed at a high frequency on human breast, colon, ovary and lung carcinomas.

Gillies, at "Magic bullets: an update on therapeutic antibodies", London 27<sup>th</sup> to 28<sup>th</sup> June 2001, reported that CT26-EpCAM subcutaneous tumours treated with <sup>125</sup>I-KS-IL2 (iodine-125 labelled KS antibody IL-2 fusion) and Taxol® showed the two treatments to have a synergistic effect when the immunotherapy was given 24 hours after the chemotherapy. From this he concluded that "Optimal chemotherapeutic doses may lower tumour interstitial pressure and increase targeting of immunocytokines", in other words the chemotherapy e.g. Taxol®, should be administered before the immunotherapeutic.

10

Thus, both DeNardo and Gillies suggest that the order of administration is important to the efficacy of a combined chemotherapy and immunoradiotherapy. However they disagree on the most useful order of administration for the chemotherapeutic and immunotherapeutic agents.

15

The search for anti-cancer agents and methods of treatment is ongoing and intense. The present invention seeks to provide further agents and methods for the treatment of cancers.

## 20 Summary of the invention

The inventor has discovered that a synergistic tumouricidal effect can be obtained by means of a combined treatment with a radiolabelled antibody that binds selectively to polymorphic epithelial mucin (PEM), and a chemotherapeutic agent.

PEM is a component of the human milk fat globule. PEM is expressed by cells in several body tissues and is also found in urine. Significantly, PEM is known to be expressed in epithelial cancer cells, notably in ovarian, gastric, colorectal and pancreatic cancer cells.

30

The preferred chemotherapeutic agent is the antineoplastic agent Taxotere® (Docetaxel), which is a semi-synthetic analogue of Taxol®. For an overview of Taxotere® see J L Fabre *et al.* (1995) *Drugs Future*, 20, pp. 464-471. For synthesis and structure see M Colin *et al.* US 4924012 (to Rhône-Poulenc Sante). For anti cancer activity see Riou *et al.* (1992) *Biochem. Biophys Res. Commun.* 187, pp. 164-170. Taxotere® is available commercially from Rhône-Poulenc Rorer. Other preferred chemotherapeutic agents are: Cisplatin (Faulding), Cyclophosphamide (Pharmacia and Upjohn), Vincristine (Faulding) and Gemcitabine (Lilly).

10

Monoclonal antibodies that will bind to PEM are already known, but in any case, with today's techniques in relation to monoclonal antibody technology, antibodies can be prepared to most antigens. Suitable monoclonal antibodies to selected antigens may be prepared by known techniques, for example those disclosed in "*Monoclonal Antibodies: A manual of techniques*", H Zola (CRC Press, 1988) and in "*Monoclonal Hybridoma Antibodies: Techniques and Applications*", J G R Hurrell (CRC Press, 1982) and "*Antibody Engineering, A Practical Approach*", J McCafferty *et al.*, ed. (IRL Press, 1996).

20

WO 01/74905 discloses antibodies that bind selectively to PEM and are useful in accordance with the present invention.

Preferably, the antibody is HMFG-1, which is available from Imperial Cancer Research Fund, England. More preferably the antibody is a humanised HMFG-1. Such antibodies are disclosed in WO 92/04380.

HMFG antibodies are raised against human milk fat globule (HMFG), in a delipidated state (see Taylor-Papadimitriou *et al.*, (1981), *Int. J. Cancer* 28 pp. 17-21 and Gendler *et al.*, (1988), *J. Biol. Chem.* 236 pp. 12820-12823).

30

HMFG-1 monoclonal antibodies bind to a particular component of HMFG, namely polymorphic epithelial mucin (PEM). Binding is thought to involve the amino acid sequence APDTR within the twenty amino acid tandem repeats of the *muc-1* gene product.

By 'humanised antibody' we include monoclonal antibodies having at least one chain wherein the framework regions are predominantly derived from a first, acceptor monoclonal antibody of human origin and at least one complementarity-determining region (CDR) is derived from a second, donor monoclonal antibody that may be of human or non-human origin, for example it may be a murine monoclonal antibody.

Preferably both chains of the humanised monoclonal antibody CDRs are grafted from a donor monoclonal antibody having specificity for PEM.

Advantageously, the CDR-grafted (*i.e.* humanised) chain comprises two or all three CDRs derived from a donor antibody having specificity for PEM.

Conveniently, the humanised monoclonal antibody comprises only human framework residues and CDRs from a donor antibody having specificity for PEM.

However, it will be appreciated by those skilled in the art that in order to maintain and optimise the specificity of the humanised antibody it may be necessary to alter one or more residues in the framework regions such that they correspond to equivalent residues in the donor antibody.

Conveniently, the framework regions of the humanised antibody are derived from a human IgG monoclonal antibody.

Methods of making humanised monoclonal antibodies are well-known in the art, for example see Jones *et al.* (1986) *Nature* 321 pp. 522-525, Riechmann *et al.* (1988) *Nature* 322 pp. 323-327, Verhoeyen *et al.* (1988) *Science* 239 pp. 1534-1536 and EP 239 400 (to Winter).

By "antibody" we include antibody fragments and antigen binding molecules. These molecules include Fab-like molecules (Better *et al.* (1988) *Science* 240, 1041); Fv molecules (Skerra *et al.* (1988) *Science* 240, 1038); single-chain Fv (ScFv) molecules where the  $V_H$  and  $V_L$  partner domains are linked via a flexible oligopeptide (Bird *et al.* (1988) *Science* 242, 423; Huston *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85, 5879) and single domain antibodies (dAbs) comprising isolated V domains (Ward *et al.* (1989) *Nature* 341, 544). A general review of the techniques involved in the synthesis of antibody fragments which retain their specific binding sites is to be found in Winter & Milstein (1991) *Nature* 349, 293-299.

By "ScFv molecules" we mean molecules wherein the  $V_H$  and  $V_L$  partner domains are linked via a flexible oligopeptide.

20

The advantages of using antibody fragments, rather than whole antibodies, are several-fold. The smaller size of the fragments may lead to improved pharmacological properties, such as better penetration of solid tissue. Effector functions of whole antibodies, such as complement binding, are removed. Fab, Fv, ScFv and dAb antibody fragments can all be expressed in and secreted from *E. coli*, thus allowing the facile production of large amounts of the said fragments.

Whole antibodies, and  $F(ab')_2$  fragments are "bivalent". By "bivalent" we mean that the said antibodies and  $F(ab')_2$  fragments have two antigen

30

combining sites. In contrast, Fab, Fv, ScFv and dAb fragments are monovalent, having only one antigen combining sites.

By "in combination with one another" regarding the antibody and  
5 chemotherapy agent treatments we include the meaning not only that the antibody and chemotherapy agents are administered simultaneously, but also that they are administered separately and sequentially.

Preferably the antibody and chemotherapy agents are  
10 administered between 0 and 24 hours apart with either the antibody or the chemotherapy being administered first.

By "binds selectively" we include the meaning that the antibodies in  
question will specifically bind cells displaying PEM on their surface and  
15 will not bind to those cells not displaying PEM.

By "treatment" we include the meanings that the tumour size is reduced and/or further tumour growth is retarded and/or prevented and/or the tumour is killed.

20

Examples embodying an aspect of the invention will now be described with reference to the following figures in which:

Figure 1 shows the effect of various treatments on tumour volume in  
25 a human derived bladder cancer cell line subcutaneously implanted on a mouse.

Figure 2 shows the effect on tumour tripling times of various treatments.



*Example 1: Combination therapy increases tumour-tripling times significantly.*

## Materials & methods

5

### *Cell lines*

The human bladder tumour cell line, HT1376, and the human colon tumour cell line HT29 expressing polymorphic epithelial mucin (PEM) were cultured in RPMI 1640 tissue culture medium containing 100 U·ml<sup>-1</sup> penicillin and 100 µg·ml<sup>-1</sup> streptomycin, supplemented with 10% foetal calf serum in a humidified atmosphere of 5% carbon dioxide in air. HT1376 is a human bladder carcinoma cell line obtained from the European Collection of Animal Cell Cultures (ECACC no. 87032402). HT29 is human colon carcinoma cell line obtained from European Collection of Animal Cell Cultures (ECACC no. 91072201)

15

### *Antibody*

The fully humanised version of the anti-PEM antibody, HMFG1, was produced by Lonza, Slough, UK. This humanised HMFG1 (hHMFG1) was conjugated with the chelating agent CITC-DPTA by BioInvent, Sweden.

20

### *Radiolabelling*

CITC-DTPA-conjugated hHMFG1 was radiolabelled with <sup>90</sup>Y in acetate buffer (pH 5.5) at room temperature for 30 minutes. Disodium EDTA was added to the reaction mixture such that the final EDTA concentration was 5 mM and left to stand at room temperature for approximately 10 min. The radiolabelled protein was then purified by size exclusion chromatography and the protein-containing fractions pooled.

25

### *Animal model*

#### *Mice*

Female MF1 athymic nude (nu/nu) mice were used throughout these studies. The mice were bred at the Biological Research Facility of St. George's Hospital Medical School and were housed in sterile filter cages and maintained on irradiated diet and sterile water. Tumours were established by subcutaneous injection of  $5 \times 10^6$  cells in the right flank.

#### *Combination therapy*

Stock solutions of drugs: Taxotere (Aventis), Cisplatin (Faulding), Cyclophosphamide (Pharmacia and Upjohn), Vincristine (Faulding) and Gemcitabine (Lilly) were diluted in saline and injected intravenously into tumour bearing nude mice via a lateral tail vein. Humanised HMFG1 anti-PEM antibody (previously conjugated with the chelating agent CITC-DTPA) was radiolabelled with yttrium-90 ( $^{90}\text{Y}$ ) to a specific activity of approximately 1-2 MBq per 10  $\mu\text{g}$ . Mice received approximately 10  $\mu\text{g}$  of protein by intravenous injection.

For combination therapy, the drug and radioimmunotherapy (RIT) were given by sequential injections into the two lateral tail veins. The drug was given either 24 h before or 24 h after the RIT. In the case of Cisplatin, this was also given at the same time as the RIT. Control mice were untreated.

#### *Tumour therapy schedule*

Approximately three weeks after tumour inoculation, when the tumours were around  $0.2 \text{ cm}^3$  in volume (7-8 mm in diameter), mice were

divided into treatment groups of 7-8 mice each. The tumour volumes in each group at the time of treatment were not significantly different. Mice were injected with various doses of  $^{90}\text{Y}$ -labelled hHMFG1 radioimmunotherapy (10-20  $\mu\text{g}$ , 1.2-2.0 MBq) either alone or in  
5 combination with the test drug given either 24 h before or 24 h after radioimmunotherapy, or with chemotherapy or chemotherapy vehicle alone. One group of mice was left untreated as a control on some occasions when experimental animals were treated. Tumour diameters ( $d_1$ ,  $d_2$  and  $d_3$ ) were measured twice weekly in three orthogonal directions using a vernier  
10 calliper and the tumour volume ( $v$ ) calculated according to the formula for an ellipsoid:

$$v = \frac{\pi}{6} (d_1 \cdot d_2 \cdot d_3)$$

Tumour measurement commenced one week before treatment and continued until the tumours had at least tripled in volume. Relative tumour volume (the volume of each tumour divided by the tumour volume on the day  
15 of treatment) was calculated to minimise the effect of variation in treatment volume of the individual tumours. The end-point was defined as time for the relative tumour volume to reach 3.

### *Statistics*

20 The Wilcoxon rank sum test was used to compare the groups of mice receiving the various treatment protocols. A  $p$ -value  $< 0.05$  was considered to be significant.

**Results**

Volume tripling times (days)  
<sup>90</sup>Y-hHMFG1 plus chemotherapy

5 **Table 1**

Treatment	Median	T-C
Control (18-19)	17.9	
Taxotere® 10mg/kg (15)	43.8	25.9
1.2 MBq <sup>90</sup> Y-hHMFG1	23.4	5.5
2.0 MBq <sup>90</sup> Y-hHMFG1	46.5	28.6
1.2 MBq + Taxotere®	73.3	55.4
Control (25-26)	19.0	
Taxotere® 10 mg/kg (27-31)	41.2	22.2
Dox 10 mg/kg (20-22)	52.9	33.9
Taxol® 10 mg/kg (23)	30.2	11.2
Control (76-79)	21.7	
1.6 MBq <sup>90</sup> Y-hHMFG1	40.6	18.9
Tax + dox 5 mg/kg each	35.9	14.2
1.6 MBq + chemo	49.8	28.1
Chemo + 1.6 MBq (80)	50.8	29.1

Table 2HT1376 Taxotere

Treatment	Median	T-C
Observer 1		
Control	21.9	
1.2 MBq <sup>90</sup> Y-hHMFG1	23.8	1.9
10 mg/kg Taxotere	43.5	21.6
Taxotere + 1.2 MBq	49.5	27.6
1.2 MBq + Taxotere	51.7	29.8

*HT1376 Cisplatin*

Treatment	Median	T-C
Observer 1		
Control	21.9	
1.2 MBq <sup>90</sup> Y-hHMFG1	22.9	1.0
10 mg/kg Cisplatin	28.8	6.9
Cisplatin + 1.2 MBq (-24 h)	65.9	44.0
10 mg/kg + 1.2 MBq (0 h)	58.4	36.5
1.2 MBq + Cisplatin (+24 h)	27.4	15.5
Control	21.9	
1.4 MBq <sup>90</sup> Y-hHMFG1	30.9	9.0
10 mg/kg Cisplatin	28.8	6.9
10 mg/kg + 1.4 MBq (-24 h)	46.6	24.7
Control	21.9	
1.4 MBq <sup>90</sup> Y-hHMFG1	30.9	9.0
2 mg/kg Cisplatin	28.9	7.0
2 mg/kg + 1.4 MBq (0 h)	37.5	15.6

*HT29 gemcitabine*

Treatment	Median	T-C
Observer 1		
Control	8.8	
1.4 MBq <sup>90</sup> Y-hHMFG1	15.7	6.9
240 mg/kg gemcitabine	15.4	6.6
Gemcitabine + 1.4 MBq	15.8	7.0
1.4 MBq + gemcitabine	19.0	10.2
Observer 2		
Control	9.5	
1.4 MBq <sup>90</sup> Y-hHMFG1	14.4	4.9
240 mg/kg gemcitabine	12.8	3.3
Gemcitabine + 1.4 MBq	14.6	5.1
1.4 MBq + gemcitabine	12.2	2.7

*HT29 cyclophosphamide*

Treatment	Median	T-C
<b>Observer 1</b>		
Control	10.2	
1.4 MBq <sup>90</sup> Y-hHMFG1	11.8	1.6
200 mg/kg cyclo	15.5	5.3
Cyclo + 1.4 MBq	13.0	2.8
1.4 MBq + cyclo	17.4	7.2
<b>Observer 2</b>		
Control	10.7	
1.4 MBq <sup>90</sup> Y-hHMFG1	13.2	2.5
200 mg/kg cyclo	14.1	3.4
Cyclo + 1.4 MBq	13.8	3.1
1.4 MBq + cyclo	17.5	6.8



*HT29 vincristine*

Treatment	Median	T-C
<b>Observer 1</b>		
Control	10.2	
1.4 MBq <sup>90</sup> Y-hHMFG1	11.8	1.6
2 mg/kg vincristine	13.4	3.2
Vincristine + 1.4 MBq	15.6	5.4
1.4 MBq + vincristine	16.4	6.2
<b>Observer 2</b>		
Control	10.7	
1.4 MBq <sup>90</sup> Y-hHMFG1	13.2	2.5
2 mg/kg vincristine	14.7	4.0
Vincristine + 1.4 MBq	14.2	3.5
1.4 MBq + vincristine	16.6	5.9

The tables summarise the treatments given and the results obtained and show the median tumour volume tripling times for HT1376 bladder tumour xenografts treated with Taxotere or Cisplatin and for HT29 colon tumour xenografts treated with Gemcitabine, Cyclophosphamide, or Vincristine. The second column is the treated minus control tripling time, i.e. the advantage of the drug or drug combination over untreated tumours. Hence, the larger the T-C value, the better the therapeutic effect obtained. Two sets of data are presented, due to tumour measurements being taken by two observers. Figures 1 & 2 represent graphically a selection of the results obtained.

From Table 1 it is clear that a combination of 1.2 MBq <sup>90</sup>Y-hHMFG1 and Taxotere® increased the tumour tripling time more than twice as much as Taxotere® on its own and ten times as much as the same dose of <sup>90</sup>Y-

hHMFG1 on its own. In fact, the contribution of radiolabelled antibody and Taxotere® was almost twice as effective ( $T-C = 55.4$ ) as the additive effect of the individual treatments ( $T-C = 5.5 + 25.9 = 31.4$ ).

5 Table 1 also shows that the sequence of treatment with  $^{90}\text{Y}$ -hHMFG1 and Taxotere® does not alter the effectiveness of the treatment significantly. Administration of 1.6MBq  $^{90}\text{Y}$ -hHMFG1 before Taxotere® gave a T-C value of 28.1 whereas administration in the reverse order gave a T-C value of 29.1.

10

From Table 2 it is clear that for example a combination of 1.2 MBq  $^{90}\text{Y}$ -hHMFG1 and Taxotere® increased the tumour tripling time more than 50% over Taxotere® on its own and fifteen times as much as the same dose of  $^{90}\text{Y}$ -hHMFG1 on its own. The contribution of radiolabelled antibody and Taxotere® was 25% more than the additive effect of the individual treatments ( $T-C = 29.8$  in combination,  $T-C = 1.9 + 21.6 = 23.5$ ).

15 Table 2 also shows that the sequence of treatment with  $^{90}\text{Y}$ -hHMFG1 and Taxotere® does not alter the effectiveness of the treatment significantly. Administration of 1.2MBq  $^{90}\text{Y}$ -hHMFG1 before Taxotere® gave a T-C value of 29.8 whereas administration in the reverse order gave a T-C value of 27.6.

20 Table 2 also shows that Cisplatin demonstrates a synergistic effect no matter the order of administering the radioimmunotherapy and the drug. For example, the 1.2 MBq dose gives a T-C value of 44, 36.5 and 15.5 over the different dosage regimes, the smallest of which is double the combined addition of the drugs administered individually at a T-C value of 7.9. The most effective treatment was to administer the drug 24 hours before the radioimmunotherapy. Gemcitabine also showed an overall increase in T-C values when the drug and radioimmunotherapy were given in combination.

25

30

Cyclophosphamide and vincristine were both most effective when given after the radioimmunotherapy. In that treatment they showed a synergistic effect over the mere additive effect of the drugs administered individually. Vincristine is also effective when administered before the radioimmunotherapy although to a slightly lesser extent.

Overall, the data shows that the combination therapy is more effective than either treatment administered alone. The effects are of differing amounts but represent a synergistic effect (i.e. more than additive).

10

*Example 2: Use of combination therapy in tumour treatment*

The combination therapy experimentally tested on mouse tumours in example 1 can be applied to use in the treatment of human tumours.

15

Treatment of human tumours requires administration of the standard clinical chemotherapy dose in  $\text{mg}/\text{m}^2$  ( $\text{mg}/\text{m}^2$  is calculated approximately by multiplying  $\text{mg}/\text{kg}$  by 230) for the chemotherapeutic agent being used. The standard clinical dose for a particular patient can easily be calculated based on that patient's specific circumstances and would form part of the day to day activities of the skilled person.

The radiolabelled antibody would preferably be administered at an initially low dose e.g. 37 to 185 Mbeq (1 to 5 milliCuries) of radiation. It is envisaged that this initially low dose of radioimmunotherapy can be raised in subsequent doses, dependent on the individual requirements of the patient. Higher doses are envisaged to be administered in amounts such that up to 400 Mbeq of radioimmunotherapy can safely reach the target tumour.

25  
30

Maraveyas *et al.* (1995) Cancer Research; 55; pp 1020-102 (incorporated herein by reference), discloses that 30% of intraperitoneal

radiolabelled antibody reaches the target tumour site. Hence, it is well within the knowledge of a skilled person to apply the teaching of Maraveyas to tailoring the most appropriate radioimmunotherapy dose, based on the amount of radiation to reach the target tumour and the weight (and/or surface area) of the patient, in conjunction with any other relevant factors.

The time between administration of the chemotherapeutic agent and the radioimmunotherapy is preferably between 0 and 24 hours, with either the chemotherapeutic or the radioimmunotherapy being administered first. It is well within the skilled person's capabilities to construct a schedule of times for administering the chemotherapeutic and radioimmunotherapy based on the needs of the patient and availability of appropriate resources.

The combination therapy will be administered in a course of treatment. The exact frequency of treatment administration within the course and length of the course as a whole will depend upon the particular chemotherapeutic agent being used and the circumstances of the individual patient. It is entirely within the scope of a skilled person's abilities to be able to determine the appropriate length and frequency of treatment.

## CLAIMS

1. A therapeutic system for the treatment of tumours comprising a combination of component (i) a radiolabelled antibody and component (ii) a chemotherapeutic agent, wherein the antibody binds selectively to polymorphic epithelial mucin (PEM); the components (i) and (ii) being provided for use in the treatment of a tumour wherein the radiolabelled antibody and a chemotherapeutic agent are administered in combination with one another.
2. A therapeutic system for use as claimed in Claim 1 wherein the antibody treatment precedes treatment with the chemotherapeutic agent.
3. A therapeutic system for use as claimed in Claim 1 wherein the chemotherapeutic agent treatment precedes treatment with the antibody.
4. A therapeutic system as claimed in any of Claims 1 to 3 wherein the antibody is humanised.
5. A therapeutic system as claimed in Claim 4 wherein the antibody is HMFG-1.
6. A therapeutic system as claimed in any preceding claim wherein the chemotherapeutic agent is selected from at least one of docetaxel, paclitaxel, doxorubicin and cisplatin.
7. A therapeutic system for use as claimed in Claim 6 wherein the chemotherapeutic agent is docetaxel.

8. A therapeutic system as claimed in any of claims 1 to 5 wherein the chemotherapeutic agent is selected from at least one of gemcitabine, cyclophosphamide and vincristine.
- 5
9. A therapeutic system as claimed in any previous claim wherein the tumour is associated with at least one of the following disorders: breast cancer, ovarian cancer, lung cancer, gastric cancer, bladder cancer and squamous cell carcinoma, such as head and neck cancer.
- 10
10. A method of treating a tumour comprising exposing the tumour to a combination of component (i) a radiolabelled antibody and component (ii) a chemotherapeutic agent, wherein the antibody binds selectively to polymorphic epithelial mucin (PEM).
- 15
11. A method as claimed in Claim 10 wherein the antibody treatment precedes treatment with the chemotherapeutic agent.
12. A method as claimed in Claim 10 wherein the chemotherapeutic agent treatment precedes treatment with the antibody.
- 20
13. A method as claimed in any of Claims 10 to 12 wherein the antibody is humanised.
- 25
14. A method as claimed in Claim 13 wherein the antibody is HMFG-1.
15. A method as claimed in any of claims 10 to 14 wherein the chemotherapeutic agent is selected from at least one of docetaxel, paclitaxel, doxorubicin and cisplatin.

16. A method as claimed in Claim 15 wherein the chemotherapeutic agent is docetaxel.
17. A method as claimed in any of claims 10 to 14 wherein the  
5 chemotherapeutic agent is selected from at least one of gemcitabine, cyclophosphamide and vincristine.
18. A method as claimed in any of claims 10 to 17 wherein the tumour is  
10 associated with at least one of the following disorders: breast cancer, ovarian cancer, lung cancer, gastric cancer, bladder cancer and squamous cell carcinoma, such as head and neck cancer.
19. Use of a combination of component (i) a radiolabelled antibody and  
15 component (ii) a chemotherapeutic agent, wherein the antibody binds selectively to polymorphic epithelial mucin (PEM) in the manufacture of a therapeutic system for the treatment of cancer.
20. A use as claimed in Claim 19 wherein the antibody is humanised.
- 20 21. A use as claimed in Claim 20 wherein the antibody is HMFG-1.
22. A use as claimed in any of claims 19 to 21 wherein the  
25 chemotherapeutic agent is selected from at least one of docetaxel, paclitaxel, doxorubicin and cisplatin.
23. A use as claimed in Claim 22 wherein the chemotherapeutic agent is  
docetaxel.
24. A use as claimed in any of claims 19 to 21 wherein the  
30 chemotherapeutic agent is selected from at least one of gemcitabine, cyclophosphamide and vincristine.

25. A use as claimed in any of claims 19 to 24 wherein the tumour is associated with at least one of the following disorders: breast cancer, ovarian cancer, lung cancer, gastric cancer, bladder cancer and squamous cell carcinoma, such as head and neck cancer.
- 5
26. A therapeutic system substantially as described herein with reference to one or more of the examples.
- 10
27. A method of treatment substantially as described herein with reference to one or more examples.
28. Use of a combination of component (i) a radiolabelled antibody and component (ii) a chemotherapeutic agent, wherein the antibody binds selectively to polymorphic epithelial mucin (PEM) in the manufacture of a therapeutic system for the treatment of cancer substantially as described herein with reference to one or more examples.
- 15




$^{90}\text{Y}$ -hHMFG1 plus chemotherapy

Figure 1

CONFIDENTIAL 12 JUL 2004

✓

**THIS PAGE BLANK** (USPTO)

## HT1376 combination therapy




Figure 2

DT04 Rec'd PCT 1 2 JUL 2004

**THIS PAGE BLANK (USPTO)**

# INTERNATIONAL SEARCH REPORT

In tional Application No  
PCT/GB 03/00105

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 A61K39/395

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)  
EPO-Internal, WPI Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00 61185 A (BELLET ROBERT E ; VOGEL CHARLES L (US)) 19 October 2000 (2000-10-19) ---	
A	WO 01 74905 A (ANTISOMA RES LTD ; YOUNG ROBERT JAMES (GB)) 11 October 2001 (2001-10-11) cited in the application ---	
A	WO 92 04380 A (UNILEVER PLC ; UNILEVER NV (NL)) 19 March 1992 (1992-03-19) cited in the application ---	
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

15 April 2003

Date of mailing of the international search report

06/05/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Bardili, W

# INTERNATIONAL SEARCH REPORT

Int'l Application No  
PCT/GB 03/00105

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DENARDO, S.J. ET AL.: "Synergy of taxol and radioimmunotherapy with yttrium-90-labeled chimeric L6 antibody: Efficacy and toxicity in breast cancer xenografts"</p> <p>PROC. NATL. ACAD. SCI. USA, vol. 94, 1997, pages 4000-4004, XP002238438</p> <p>cited in the application</p> <p>-----</p>	

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/GB 03/00105

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0061185	A	19-10-2000	AU 4332700 A	14-11-2000
			BR 0009647 A	08-01-2002
			CN 1350464 T	22-05-2002
			CZ 20013614 A3	17-07-2002
			EP 1169059 A1	09-01-2002
			HU 0201459 A2	28-09-2002
			NO 20014842 A	04-10-2001
			TR 200102845 T2	21-01-2002
			WO 0061185 A1	19-10-2000
			US 6333348 B1	25-12-2001
WO 0174905	A	11-10-2001	AU 4430701 A	15-10-2001
			WO 0174905 A1	11-10-2001
			US 2002122798 A1	05-09-2002
WO 9204380	A	19-03-1992	AT 233279 T	15-03-2003
			AU 653167 B2	22-09-1994
			AU 8495391 A	30-03-1992
			BG 60716 B1	31-01-1996
			BG 97607 A	31-03-1994
			CA 2090961 A1	08-03-1992
			DE 69133200 D1	03-04-2003
			EP 0483961 A1	06-05-1992
			FI 930984 A	05-03-1993
			WO 9204380 A1	19-03-1992
			HU 67796 A2	28-04-1995
			JP 6500468 T	20-01-1994
			JP 2003061689 A	04-03-2003
			NO 930825 A	05-05-1993
			NO 20014822 A	05-05-1993
			RO 113432 B1	30-07-1998
			US 2002086978 A1	04-07-2002
			US 6204366 B1	20-03-2001

**THIS PAGE BLANK** (USPTO)



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO**